



International Journal of Surgery Science

E-ISSN: 2616-3470
P-ISSN: 2616-3462
© Surgery Science
www.surgeryscience.com
2020; 4(1): 354-356
Received: 18-11-2019
Accepted: 21-12-2019

Dr. Entesar Mahmood Amer
AL- Najaf Health Directorate, Iraq

Dr. Ahlam Mahdi Mousa
AL- Najaf Health Directorate, Iraq

Membranous rupture in relation to c-reactive protein

Dr. Entesar Mahmood Amer and Dr. Ahlam Mahdi Mousa

DOI: <https://doi.org/10.33545/surgery.2020.v4.i1.g.361>

Abstract

The aim of the study was to prediction of sub rupture of membranes (PPROM) and preterm labour. Maternal and cord blood for C (CRP) sensitivity as a confirmative test for sub with PPRM as study group and 30 pregnant of same gestat in labour. All submitted to CRP test, high vaginal swab, placental swab for C/S and cord blood for CRP. CRP positive in study group 18 cases (60%) and control group 8 cases (36.6%) for urine C/S in study group was 11 positive in study group 8 cases (27.3%) and in control group 3 cases (10%), CRP test, urine for C/S and high the vag. for C/S were statistically not significant CRP test sensitivity was 78% and specificity was 62.5% So CRP test can used as screening test for infection.

Introduction: Preterm pre labour rupture of membranes (PPROM) mean rupture of fetal membranes after 24 weeks gestation and before 37 completed weeks or 259 completed days of gestation. Preterm birth: mean birth occurs before 37 completed weeks or 259 completed days of gestation. It is a leading cause of infant death and several contributing mechanisms to this morbidity have been identified over the past 10 years Macdonald Gant, 2000). It is obvious that seve involved in the pathogenesis of preterm birth, which may explain why it has proved so difficult to predict and prevent, the too early activation of the fetal hypothalamic adrenal axis may result from maternal psychosocial or fet physiological stress. Such physiological fetal stress may in turn be a consequence of microbial invasion of fetal membranes, amniotic fluid and the fetus itself. The critical mediator of stress-induced preterm birth appear to be corticotrophin releasing hormone. It stimulates the production of prostaglandins by cells in the amnion, placenta, chorion and urine deciduas (Staffan Bergstrom, 2003). membranes "chorioamnionitis" is a cause of preterm labour in 10-20% of cases with intact membranes, this figure is higher if labour fallows preterm pre labour rupture of Membranes.

Pathophysiology: Preterm parturition due to infection is initiated by security products resulting from monocyte activation cytokines including interleukin I.

Keywords: Infection, CRP, Screening

Introduction

sub-clinical infection as a cause of preterm estimation as screening test and urine, high vaginal swab and placental swabs for culture and sub-clinical infection. Thirty pregnant of (24 gestation age with intact fetal membranes but cases (36.6%) and in control group 2 cases (6.6%) for high the vaginal swab swab for C/S all were statistically significant. Cord blood C is an open access article distributed under the Creative Commons Att provided the original work is properly cited. entify (Cunningham). Several pathways are hypothalamic- pituitary adrenal fetal. Corticotrophin the). Local infection of branes, Tumour necrosis factors (TNF) and interleukin implicated in preterm labour. TNF and matrix. metalloproteinase also promote programmed death of amniotic cells, the combined effect of these mechanisms may provoke preterm birth (Lok Wood and Kuczynski, 1999 [4] access for bacteria with intact membranes is unclear but Escherichia coli can permeate living chorio membranes thus intact fetal membranes at the cervix are not necessarily a barrier to ascending bacterial I amniotic fluid and figure (Cunningham Macdonald Gant, 2000 [1] shows pathway for bacterial initiation of pretenn labour which may not required colonization of the amniotic fluid e.g. the cytokine net work of cell mediated immunity can be active locally in decidual tissue that line the fore bag fetal membranes

(Cunningham Macdonald Gant, 2000 [1] have shown a correlation between preterm labour and asymptomatic bacteria and lower genital tract infection reactive Oxygen species which are generated by the body's response to diverse insults such as infection have also attracted attention, such insult activate collagenolytic enzymes and impair fetal membrane integrity and this impairment is then inhibited by antioxidants like vit. E 2001).

Corresponding Author:
Dr. Entesar Mahmood Amer
AL- Najaf Health Directorate, Iraq

Other Causes of PPROM

1. Placental abruption.
2. Dynamic uterine dysfunction, uterine atony or uterine inertia. preterm- pre labour C-reactive protein 24-37 weeks gestation ion CRP and placental swab Attribution License, which permits interleukin-6 are such secretory products 1999). The rout of chorio-amniotic invasion of the 2000) activated 2000). A number of studies infection species e and vit. C. (Woods *et al.*,)
3. Anemia. Recent evidence indicates that signs of inflammation or infection are prevalent in women with anemia. Ther is increased CRP concentration in more than 50% of anemic women means anemia is a sign of maternal morbidity indicating inflammation or infection of unknown origin (Svigo *et al.*, 1999) [10]. Recent literature shows that detection and estimation of surrogate marker such as CRP (C-reactive protein) cytokines and fetal fibronectin help in diagnosing intramniotic infection and in predicting and diagnosing early onset neonatal infections (Lu *et al.*, 2000) [5].

Diagnosis of PPROM

1. History: Women feeling agush of fluid vaginally.
2. A sterile speculum examination to confirm the diagnosis if no liquor is apparent.
3. Reduce fetal movement
4. Ultra-sound should not used as primary means of diagnosis of

PPROM

Laboratory Test for Diagnosis of PPROM:

1. Notarize test for vaginal PH.
2. Fern test: result from drying out of salts containing amniotic fluid.
3. Intra amniotic fluorescein and amnioscopy invasive method, and may enhance the infection.
4. Fetal fibronectin it confirm the rupture membranes and predicted for preterm labour. But it increase perinatal infection (Svigo, 1999) [10].

Complication of PPROM

1. Maternal complication:
 - a. Chorioamnionitis.
 - b. Endomyometritis.
 - c. Abruptio placentae.
 - d. Psycho-social sequelae particularly maternal hospitalization,
 - e. Induction of labour may lead to prolonged latent stage 16-20 hours which increased the likelihood of operative delivery (Svigo, 1999) [10].
2. Fetal complication:
 - a. Fetal pulmonary hypoplasia,
 - b. Skeletal deformities "talipes".
 - c. Amniotic adhesion and bands of fibrous tissue between the fetus and amnion which may cause auto-amputation,
 - d. Cord compression or prolaps, e- Fetal infection and neonatal sepsis.
 - e. Fetal intrapartum hypoxia and birth trauma associated with preterm labour involving the very low birth weight infant, whether birth is by the vaginal or abdominal rout with contribute to the perinatal risk (Svigo, 1999) [10].

C-Reactive Protein "CRP": The CRP is a plasma protein molecule consisting of a pentamer of non-glycosylated

polypeptide sub-unit that is produced by hepatocytes. This plasma protein normally is found in plasma of healthy persons in trace amounts "< 1 mg/dl" within hours of an acute injury as well as the onset of most types of inflammation or infection, the rate of production of CRP increases markedly with up to 3,000 fold increases in plasma concentration. For this reason, CRP is considered an "acute phase" protein, the plasma level of CRP rapidly decreases toward the normal range (Narinder, 1997) [7], and CRP increased and decreased more quickly than the red sedimentation rate (ESR), and the value of the CRP in the potentially infected patient exceeds Leukocyte indices. CRP production is stimulated by inter leukin 6 (IL-6), therefore IL-6 is an early and sensitive marker of infection or inflammation, so that the IL-6 should rise before CRP level rise. Median CRP values during pregnancy are higher than values for non pregnant individuals. These values are elevated further in labour. In woman not in labour 95% of values were 1.5 mg/dl or less and gestational age didn't affect serum level. In true infection the test for CRP become positive after 12 hours, so estimation of CRP at presentation may not be much value in prediction. So serial determination may be required and have better predictive value than single estimation (Staffan Bergstrom, 2003) [9]. Mild CRP elevation "within normal, non acute-phase rang" is emerging as a valuable marker of cardiovascular risk (Jam Coli, 2003) [3].

Method of GRP Test: C-reactive protein test "slide agglutination method". In this test using latex reagent which is a polystyrene latex particles of uniform size coated with the IgG fraction of an anti-human CRP specific serum. Visual observation of Ag-Ab reaction "CRP-IgG anti CRP" if reaction takes place due to presence of CRP in the serum a clear agglutination become evident. A clear agglutination will appear if the serum contains more than 6 mg/L of CRP and if no agglutination mean serum contains less than 6 mg/L of CRP which is normal range.

Reagents

- a. Latex reagent: Suspension of polystyrene latex particles coated with IgG anti-CRP in a buffer.
- b. Positive control: Diluted human serum containing more than 10 mg/L of CRP ready to use.
- c. Negative control: Diluted human serum containing less than 1 mg/L of CRP. All latex reagent positive, control, and negative contain <0.1% Sodium azid. C-reactive protein assay by using Laser nephelometry measurement of serum CRP concentrations by rate nephelometry using a Beckman Array system protein analyzer, "which is not available here" A double antibody sandwich enzyme immunoassay for C.R.P. which is high sensitivity CRP test (Saunders, 2003) [8]. Aim of the Study: To evaluation C-reactive protein as a predictor of sub-clinical infection in preterm-prelabour rupture of fetal membranes and in preterm labour with intact fetal membranes.

Patient and Methods: Present study was carried out in Maternity and Childhood Teaching Hospital in Najaf, from June to October during a year 2003. Selection of Patient: The study consist of 30 cases selected with PPROM beyond 24 weeks of gestation up to 37 incomplete weeks of gestation, "case group". A second group "control group", 30 cases without rupture of fetal membranes but in labour, with same gestational age. Both groups were subjected to various investigation "urine and high vaginal swab for culture and sensitivity and placental swab for culture and sensitivity" as a confirmative tests for sub-clinical

infection, patient who are excluded from study were:

1. Evidence of systemic infection.
 2. Mai presentation.
 3. Twin pregnancy.
 4. Polyhydramnios.
 5. Connective tissue disease. Investigation: L Evaluation of CRP by slide agglutination method of mother at admission, as mentioned above.
1. High vaginal swab for culture and sensitivity.
 2. Urine for culture and sensitivity.
 3. Cord blood for CRP estimation.
 4. Placental swab for culture and sensitivity. Parameter of Infection after delivery The mother was followed after delivery for evidence of:
 1. Maternal fever during 24 hours after delivery.
 2. Un healthy lochia "offensive discharge". Neonatal requiring for neonatal intensive care unit admission. Each new born was admitted after delivery immediately to NICU for observation apgar scor for the first and five minutes of delivery and follow up of the baby during period of admission.

Statistical Analysis: Using Chi-square test "x²". The P value for positive cases <0.05 which mean statistically significant.

Results

In this study 30 patients with PPRM as study cases and 30 patents with preterm labour and intact fetal membranes, as control cases. Both groups, Maternal and cord blood CRP estimation as a screening test for sub-clinical infection as a cause of PPRM and preterm labour. Both mother and neonates were observed post delivery for evidence of infection like maternal fever "unhealthy lochia" and neonate admission in NICU.

Urine culture

- Six cases isolated M.O were E-coli.
- Two cases the M.O were staphylococcus.
- Three cases klebsella.

Vaginal swab cultures isolate

- One cases enterobacter.
- Four cases E-coli.
- Two cases staphylococcus aureus.
- One case klebsella.
- .Three cases monillial infection

Discussion

The hypothesis that infection is the cause of PPRM stands proved. Levels of CRP rise when there is a microbial infection or an inflammation without microbes. This is supported by obtaining statistically significant maternal CRP values in patients with PPRM, which is good predictor of sub-clinical infection specially those patients who were PPRM and not in labour.

- In this study the sensitivity of CRP is 78% and specificity is 62.5% while B.R daial got 79% sensitivity and 80% specificity in predictive effectiveness of CRP as a marker of infection (Desai *et al.*, 1997) [2].
- The difference in the specificity of the test may be due to using another confirmative method such as placental and fetal membranes histopathological study for presence of evidence of inflammation or infection. And may be we have no culture media or absence of other media for vaginal infection.

- Melee holds that urinary tract infection is the likely focus predisposing to PPRM (Malee, 1992) [6] while in this study the urinary tract infection and lower genital tract infection have correlation with PPRM as demonstrated in table 2 and 3. Cervical swab in this study not performed due to deficiency in our laboratory facilities, placental swab for culture revealed 5 cases out of 30 cases showed +ve growth (16.5%) in study group and no cases in control group. And the microorganism
 - which were grown in the vaginal swabs are defer from that micro-organisms which grown in the
 - placenta or urinary tract. In the same patient, that mean there were poly microbial infection may cause the condition (Staffan Bergstrom, 2003) [9].
 - Rising of maternal CRP was not are liable predictor of prenatal neonatal infectious morbidity intensive care unit NICU this due to pre-maturity problems and not due to neonatal infection.

However maternal post-delivery infection, only 2 cases (6.6%) were developed genital tract infection this low percentage it may be due to administration of parantral antibiotics pre and post delivery as a routine role in our labour room management for each case with PPRM. Comparison with study have done in Jawaherialnehru study at 1997 they have no post delivery maternal infection (Desa *et al.*, 1997) [2].

Conclusion

The conclusion of this study is that CRP test can be used a screening test for sub-clinical infection in PPRM. CRP test has sensitively of 78% and will specificity 62.5%. This test is cheap, easy performed and available and within a short period by agglutination test which take 10 minutes in getting a results. So that, it can be used as a first line for prediction of subclinical infection for both maternal and neonatal infection.

References

1. Cunningham Macdonald Gant, Preterm Birth William's, Obstetrics, 2nd edition. 2000; 2:202-203
2. Desai BR, Shobhana S patted, Richa. A one year case control study to evaluate the incidence of infection as a cause of preterm rupture of membranes. The journal of obstet. And gyn. 1997, 590-610.
3. Jam Coli. Cardio-CRP test. Cardiol. 2003; 41(8):1358-1363.
4. Lok Wood CJ, Kuczynski E. Markers of risk preterm delivery. J. perinat Med. 1999; 27:5-20.
5. Lu GC, Golden Bery RL. Current preterm birth. Clin. Perinatal. 2000; 27:263-284.
6. Malee P, Maureen. Urinary tract infection. OBG clinics of north America, 1992; 19(2):310.
7. Narinder MK. The use of the erythrocyte sedimentation rate. The C-reactive protein assay and procalcitonin concentration in the assessment of potentially infected patient. AM. Infect. Dis., 1997; 16(3):21-24.
8. Saunders WB. The role of CRP in the evaluation and management of infants with sepsis. Advance-Neonatal care. 2003; 3(1):3-13.
9. Staffan Bergstrom. Infection-Related morbidities in the mother, fetus and neonate. The American society for nutritional sciences, 2003; 133:1656S-1660S.
10. Svigo JM, robinson JSR. Vignes Waran. Prelabour rupture of membranes. High risk, 2nd edition: 1999, 1015-1021.
11. Woods JR, Jr Plessinger MA, Miller RK. Vitamin C and E. missing links in preventing preterm prelabour rupture of membranes. AM. 5 obstet. Gynecol. 2001, 5-10