



# International Journal of Surgery Science

E-ISSN: 2616-3470

P-ISSN: 2616-3462

© Surgery Science

www.surgeryscience.com

2018; 2(2): 19-22

Received: 18-02-2018

Accepted: 23-03-2018

**Dr. G Siva Kumar**

MS(Gen Surgery),

Assistant Professor, Department of  
Surgery, Gandhi Medical College,  
Secunderabad, Telangana, India

**Dr. D Nagender Rao**

Assistant Professor,

Department of Surgery, Gandhi  
Medical College, Secunderabad,  
India

## Open VS laparoscopic appendicectomy: A comparative study

**Dr. G Siva Kumar and Dr. D Nagender Rao**

### Abstract

**Background:** Appendicectomy is the most common surgical procedure performed in emergency surgery. Because of lack of consensus about the most appropriate technique, appendicectomy is still being performed by both open (OA) and laparoscopic (LA) methods. Laparoscopic appendicectomy is likely to have less postoperative pain, early discharge, decreased wound infection, better cosmesis and also early return to routine work. The advantages of laparoscopic appendicectomy over open appendicectomy are questioned over and over.

**Objective:** The primary objective of this study was to compare the laparoscopic approach and the conventional technique in the treatment of acute appendicitis.

**Materials and Methods:** The study included 200 cases, 100 open appendicectomy cases and 100 laparoscopic appendicectomy cases, which were randomly selected and were operated in department of surgery, Gandhi Medical College, Secunderabad. The following parameters were observed during follow up in comparison between two procedures, post operative pain; duration of analgesic use; post operative complications like vomiting, ileus, abdominal abscess and wound infection. Chi-square test and student t-test were used for statistical analysis.

**Results:** Pain score was relatively high for open group when compared to laparoscopic group. Post operative complications like vomiting and ileus were lower in laparoscopic group. There is significant reduction in incidence of post operative wound infection in laparoscopic group. Laparoscopic appendicectomy was associated with a shorter hospital stay. The return to normal activity was low for laparoscopic group compared to open group. Operative time was significantly shorter in the open group.

**Conclusions:** The laparoscopic approach is a safe and efficient operative procedure in appendicectomy and it provides clinically beneficial advantages over open method.

**Keywords:** Appendicitis, appendicectomy, laparoscopic appendicectomy, open appendicectomy

### Introduction

The appendix is a blind-ended tube connected to the cecum, from which it develops in the embryo. The most common diseases of the appendix (in humans) are appendicitis and carcinoid tumors. Appendicitis is inflammation of the appendix. Symptoms commonly include right lower abdominal pain, nausea, vomiting, and decreased appetite. However, approximately 40% of people do not have these typical symptoms <sup>[1, 2]</sup>. Appendicitis is caused by a blockage of the hollow portion of the appendix. This is most commonly due to a calcified "stone" made of feces. Inflamed lymphoid tissue from a viral infection, parasites, gallstone, or tumors may also cause the blockage. This blockage leads to increased pressures in the appendix, decreased blood flow to the tissues of the appendix, and bacterial growth inside the appendix causing inflammation. The combination of inflammation, reduced blood flow to the appendix and distention of the appendix causes tissue injury and tissue death. If this process is left untreated, the appendix may burst, releasing bacteria into the abdominal cavity, leading to increased complications <sup>[3]</sup>. The surgical procedure for the removal of the appendix is called an appendicectomy. Appendicectomy can be performed through open or laparoscopic surgery. The surgical procedure for the removal of the appendix is called an appendicectomy <sup>[4, 5]</sup>. Appendicectomy can be performed through open or laparoscopic surgery. When the surgeon uses the open approach, he makes an incision in the lower right section of the abdomen. Most incisions are less than 3 in (7.6 cm) in length. The surgeon then identifies all of the organs in the abdomen and examines them for other disease or abnormalities. The appendix is located and brought up into the wounds. The surgeon separates the appendix from all the surrounding tissue and its attachment to the cecum, and then removes it. The site where the appendix was previously attached, the cecum, is closed and returned to the abdomen.

### Correspondence

**Dr. D Nagender Rao**

Assistant Professor,

Department of Surgery, Gandhi  
Medical College, Secunderabad,  
India

The muscle layers and then the skin are sewn together [6, 7, 8]. When the surgeon performs a laparoscopic appendectomy, four incisions, each about 1 in (2.5 cm) in length, are made. One incision is near the umbilicus, or navel, and one is between the umbilicus and the pubis. Two other incisions are smaller and are on the right side of the lower abdomen. The surgeon then passes a camera and special instruments through these incisions. With the aid of this equipment, the surgeon visually examines the abdominal organs and identifies the appendix. The appendix is then freed from all of its attachments and removed. The place where the appendix was formerly attached, the cecum, is stitched. The appendix is removed through one of the incisions. The instruments are removed and then all of the incisions are closed [6, 7, 8]. The primary objective of this study was to compare the laparoscopic approach and the conventional technique in the treatment of acute appendicitis.

**Materials and methods**

**Place of Study:** Department of surgery, Gandhi Medical College, Secunderabad.

**Type of Study:** Prospective and observational study on 200 appendectomy cases.

**Inclusion Criteria:** Patients with clinical diagnosis of acute or

recurrent appendicitis with necessary investigations.

**Exclusion Criteria:** Patients who didn't give consent, children below the age of 10 years, pregnant women and cases of complicated appendicitis were excluded from this study.

**Methodology:** The following parameters were observed during follow up in comparison between two procedures, post operative pain using a visual analogue pain scale and duration of analgesic used in number of days. Post operative complications like vomiting, ileus, abdominal abscess and wound infection. Patients in both study groups were discharged as soon as possible and duration of stay after surgery and duration of analgesics used after surgery in number of days is noted. Wound infection was defined as discharge of pus that required surgical drainage. Intrabdominal abscess was defined as a fluid collection diagnosed at Ultrasonography or computed tomography which contained pus at ultrasonographically guided aspiration. Presented proforma was used to collect the relevant information.

**Statistical Methods:** Chi-square test and student t- test were used for statistical analysis.

**Observations and results**

Demographic and preoperative clinical data

**Table 1:** Demographic and Preoperative Clinical Data

Characteristic	Open Appendectomy	Laparoscopic Appendectomy
Frequency	100	100
Gender		
Male	65	45
Female	35	55
Mean Age	25.8 ± 9.1	22.3 ± 7.3
Co-morbidities		
Hypertension	5	3
Diabetes Mellitus	4	2
COPD	3	2
CAD	2	2

Table 1 depicts that in the present study 65(65%) patients of open appendectomy and 45(45%) patients of laparoscopic appendectomy were males. 35(35%) patients of open appendectomy and 55(55%) laparoscopic appendectomy

were females. The mean age of the patients in two groups was 25.8 ± 9.1 years and 22.3 ± 7.3 years, respectively.

**Duration of surgery**

**Table 2:** Duration of Surgery

	Appendectomy	
	Open	Laparoscopic
Mean Duration Of Surgery	46.3 ± 11.0	66.9 ± 22.1

Table 2 shows that open appendectomy is less time consuming than laparoscopic appendectomy.

**Post operative complications**

**Table 4:** Post Operative Complications

Complication	Appendectomy		P Value
	Open	Laparoscopic	
Vomiting	20(20%)	10(10%)	<0.05*
Abdominal Abscess	5(5%)	0(0%)	>0.05
Wound Infection	17(17%)	4(4%)	<0.05*
Ileus(Hours)	31.1± 8.1	18.3± 7.4	<0.05*

**Post operative pain score**

**Table 3:** Post Operative Pain Score

	Appendectomy		Statistical Analysis	
	Open	Laparoscopic	T Value	P Value
Pain Score	2.5 ± 0.3	1.4 ± 0.4	6.88	<0.05*

\*P value significant

\*P value significant

Table 3 depicts that in the present study average pain score was 2.5 ± 0.3 in open group as compared to 1.4 ± 0.4 in laparoscopic group with p value <0.05 which was statistically significant.

Table 4 depicts that post operative complications like vomiting and ileus were lower in laparoscopic group. There is significant reduction in incidence of post operative wound infection in laparoscopic group.

## Post operative hospital stay

**Table 5:** Post operative Stay

Hospital Stay	Appendicectomy	
	Open	Laparoscopic
1 day	0	14
2 days	0	43
3 days	13	30
4 days	68	13
5-9 days	8	0
10-15 days	11	0
P Value	<0.05*	<0.05*

\*P value significant

Table 5 shows that laparoscopic appendicectomy significantly reduced the hospital stay ( $P < 0.05$ ).

## Discussion

Acute appendicitis is the most common indication for abdominal surgery with a life-time incidence between 7 to 9 percent. Appendicectomy is one of the operations which are most commonly performed by the general surgeons. Open appendectomy (OA) has been the gold standard for the treatment of acute appendicitis. Laparoscopic appendicectomy (LA) has evolved since the first performed by a German Gynecologist Semm K. Laparoscopic appendicectomy has gained acceptance as a diagnostic and treatment method for acute appendicitis with the technological advances of the past two to three decades. Since then, this procedure has been widely used [4]. In spite of its wide acceptance, there remains a continuing controversy in the literature regarding the most appropriate way of removing the inflamed appendix. In the present study 65(65%) patients of open appendicectomy and 45(45%) patients of laparoscopic appendicectomy were males. 35(35%) patients of open appendicectomy and 55(55%) laparoscopic appendicectomy were females. The mean age of the patients in two groups was  $25.8 \pm 9.1$  years and  $22.3 \pm 7.3$  years, respectively. Table 2 shows that open appendicectomy is less time consuming than laparoscopic appendicectomy. Similar observations have also been reported by other studies [10, 11]. A meta-analysis of randomized controlled trial has been reported with outcomes for 3000 patients. The mean operating time was 18 minutes longer for laparoscopic appendicectomy [12]. A prospective randomized trial comparing laparoscopic appendicectomy with open appendicectomy was conducted in 158 patients by Hansen *et al.* [13] reported that despite of longer operating time, the advantages of make it a worthwhile alternative for patients with acute appendicitis. In the present study average pain score was  $2.5 \pm 0.3$  in open group as compared to  $1.4 \pm 0.4$  in laparoscopic group with p value  $< 0.05$  which was statistically significant. The similar studies done showed the incidence of emesis was lesser and post operative ileus lesser in laparoscopic group [14, 15]. Post operative complications like vomiting and ileus were lower in laparoscopic group. There is significant reduction in incidence of post operative wound infection in laparoscopic group. The return to normal activity was low for laparoscopic group compared to open group. Similar findings were observed in other studies also [16, 17]. Marzouk M *et al.* [18] showed laparoscopic appendicectomy significantly improved the postoperative wound infection rate. There was no wound infection in the laparoscopic group, whereas in open group the infection rate was 7.6. Laparoscopic appendectomy was associated with a shorter hospital stay. Other studies have shown similar findings [19, 20].

## Conclusions

The laparoscopic appendicectomy was better than the open appendicectomy with respect to pain score, lesser use of analgesics and post operative complications. Post operative recovery was good in respect with duration of hospital stay, return to normal work. The only drawback of laparoscopic appendicectomy was with the duration of surgery. Overall laparoscopic appendicectomy is better than open appendicectomy in selected patients with acute or recurrent appendicitis. The laparoscopic approach is a safe and efficient operative procedure in appendectomy and it provides clinically beneficial advantages over open method.

## References

1. Addiss DG, Shaffer N, Fowler BS, Tauxe R. The epidemiology of appendicitis and appendicectomy in the United States. *Am J Epidemiology*. 1990; 132:910-25.
2. Seem K. Endoscopic appendectomy. *Endoscopy*. 1983; 15:59-64.
3. Kurtz RJ, Heimann TM. Comparison of open and laparoscopic treatment of acute appendicitis. *Am J Surg*. 2001; 182:211-4.
4. Garbutt JM, Soper NJ, Shannon W, Botero A, Littenberg B. Meta-analysis of randomized controlled trials comparing laparoscopic and open appendectomy. *Surg Laparosc Endosc*. 1999; 9:17-26.
5. Biondi A, Grosso G, Mistretta A, Marventano S, Toscano C, Drago F *et al.* Laparoscopic vs. open approach for colorectal cancer: evolution over time of minimal invasive surgery. *BMC Surg*. 2013; 13(2):S12.
6. Grosso G, Biondi A, Marventano S, Mistretta A, Calabrese G, Basile F. Major postoperative complications and survival for colon cancer elderly patients. *BMC Surg*. 2012; 12(1):S20.
7. Biondi A, Grosso G, Mistretta A, Marventano S, Toscano C, Gruttadauria S *et al.* Laparoscopic-assisted versus open surgery for colorectal cancer: short-and long-term outcomes comparison. *J Laparoendosc Adv Surg Tech A*. 2013; 23:1-7.
8. Guller U, Hervey S, Purves H, Muhlbaier LH, Peterson ED, Eubanks S *et al.* Laparoscopic versus open appendectomy: outcomes Biondi *et al.* *World Journal of Emergency Surgery* (2016) 11:44 Page 5 of 6 comparison based on a large administrative database. *Ann Surg*. 2004; 239:43-52.
9. Wullstein C, Barkhausen S, Gross E. Results of laparoscopic vs. conventional appendectomy in complicated appendicitis. *Dis Colon Rectum*. 2001; 44:1700-5.
10. Fogli L, Brulatti M, Boschi S, Di Domenico M, Papa V, Patrizi P *et al.* Laparoscopic appendectomy for acute and recurrent appendicitis: retrospective analysis of a single-group 5-year experience. *J Laparoendosc Adv Surg Tech A*. 2002; 12:107-10.
11. Lin HF, Wu JM, Tseng LM, Chen KH, Huang SH, Lai IR. Laparoscopic versus open a appendectomy for perforated appendicitis. *J Gastrointest Surg*. 2006; 10:906-10.
12. Cueto J, D'Allemagne B, Vazquez-Frias JA, Gomez S, Delgado F, Trullenque L *et al.* Morbidity of laparoscopic surgery for complicated appendicitis: an international study. *Surg Endosc*. 2006; 20:717-20.
13. Towfigh S, Chen F, Mason R, Katkhouda N, Chan L, Berne T. Laparoscopic appendectomy significantly reduces length of stay for perforated appendicitis. *Surg Endosc*. 2006; 20:495-9.
14. Roviario GC, Vergani C, Varoli F, Francese M, Caminiti R,

- Maciocco M. Videolaparoscopic appendectomy: the current outlook. *Surg Endosc.* 2006; 20:1526-30.
15. Ortega AE, Hunter JG, Peters JH, Swanstrom LL, Schirmer B. A prospective, randomized comparison of laparoscopic appendectomy with open appendectomy. *Laparoscopic Appendectomy Study Group. Am J Surg.* 1995; 169:208-12.
  16. Milewczyk M, Michalik M, Ciesielski M. A prospective, randomized, unicenter study comparing laparoscopic and open treatments of acute appendicitis. *Surg Endosc.* 2003; 17:1023-8.
  17. Bresciani C, Perez RO, Habr-Gama A, Jacob CE, Ozaki A, Batagello C *et al.* Laparoscopic versus standard appendectomy outcomes and cost comparisons in the private sector. *J Gastrointest Surg.* 2005; 9:1174-80.
  18. Marzouk M, Khater M, Elsadek A, Abdelmoghny. Laparoscopic vs open appendectomy: a prospective comparative study of 227 patients. *Surg Endosc.* 2003; 17:721-4.
  19. Katkhouda N, Mason RJ, Towfigh S, Gevorgyan A, Essani R. Laparoscopic versus open appendectomy: a prospective randomized double-blind study. *Ann Surg.* 2005; 242:439-48.
  20. Ignacio RC, Burke R, Spencer D, Bissell C, Dorsainvil C, Lucha PA. Laparoscopic versus open appendectomy: what is the real difference? Results of a prospective randomized double-blinded trial. *Surg Endosc.* 2004; 18:334-7.