



E-ISSN: 2616-3470
P-ISSN: 2616-3462
© Surgery Science
www.surgeryscience.com
2021; 5(4): 127-129
Received: 05-07-2021
Accepted: 06-09-2021

Dr. Subhabrata Das
MS, MCH, Surgical Oncology,
Director Oncology Services, Pacific
Institute of Medical Sciences,
Udaipur, Rajasthan, India

Dr. Gururaj Deshpande
Assistant Professor,
Dept of Surgical oncology, Kidwai
Memorial Institute of Oncology,
VTSM Peripheral Cancer Centre,
Kalaburagi, Karnataka, India

Corresponding Author:
Dr. Gururaj Deshpande
Assistant Professor,
Dept of Surgical oncology, Kidwai
Memorial Institute of Oncology,
VTSM Peripheral Cancer Centre,
Kalaburagi, Karnataka, India

International Journal of Surgery Science

Anterior Glissoneon approach: A novel technique for liver resections and our experience

Dr. Subhabrata Das and Dr. Gururaj Deshpande

DOI: <https://doi.org/10.33545/surgery.2021.v5.i4b.771>

Abstract

Glisson's sheath was discovered by Johannis Walaeus in 1640. Francis Glisson published his book entitled *Anatomia Hepatis* in 1654 and accurately portrayed the essential anatomy of the major vessels of the liver. The Glisson's capsule wraps the portal vein, hepatic artery and bile duct and these three structures are therefore regarded as forming one morphological system, which since then has been referred to as the Glissonean system.

Keywords: Hepatectomy, Lobectomy, Segmentectomy, Hepatocellular Carcinoma, Anterior Approach

Introduction

Liver surgical techniques have advanced considerably with increased knowledge of the surgical anatomy of the liver. Lortat- Jacob and Robert and Honjo and Araki performed anatomical right hepatectomy with dissection of the hepatic artery, portal vein and bile duct in the hepatoduodenal ligament around 1950. This procedure has been referred to as a controlled hepatectomy method. By contrast, Lin and Tung and Quang initially dissected the liver parenchyma and then approached and ligated the portal pedicles intrahepatically. These two procedures of right hepatectomy were recognized as the basic anatomical liver resection techniques. The Glissonean pedicle approach at the hepatic hilus was reported by Takasaki *et al.* in 1986. Couinaud had also previously described a simplified left hepatectomy with the Glissonean pedicle approach at the hepatic hilus similar to Takasaki's procedure in 1985. The approaches of Couinaud and Takasaki to the Glissonean pedicles are extrahepatic approaches without liver dissection, and are completely different from those of Lin and Tung. Launois and Jamieson described the intrahepatic Glissonean pedicle approach to liver resection in 1992. This new Glissonean pedicle approach has provided in-depth knowledge of the surgical anatomy of the liver and has made different types of hepatectomy possible including not only hemihepatectomy but also small anatomical hepatectomies, such as sectionectomy and Couinaud's segmentectomy in a normal and cirrhotic liver.

The Glisson's Capsule and the hilar plate

Glisson's sheath was discovered by Johannis Walaeus in 1640. Francis Glisson published his book entitled *Anatomia Hepatis* in 1654 and accurately portrayed the essential anatomy of the major vessels of the liver, The Glisson's capsule wraps the portal vein, hepatic artery and bile duct and these three structures are therefore regarded as forming one morphological system, which since then has been referred to as the Glissonean system. The hepatic vein system is another system in the liver. Specifically, the Glisson's capsule connects to the capsule of the liver and the hepatoduodenal ligament. The Glisson's capsule forms a thick plate at the inferior part of the liver, which is referred to as the hilar plate. The hilar plate connects to the cystic plate, umbilical plate and Arantian plate. The hilar plate can be detached from the quadrate lobe because no branch originates from along the anterior margin or the upper surface of the hilar plate. The small caudate branches originate from along the posterior margin of the hilar plate. When the hilar plate is pulled down after detaching the liver parenchyma of the quadrate lobe, the right and left Glissonean pedicles can be easily approached. The most important point is that any variation, particularly in the arteries and bile ducts, occurs under the hilar plate. Therefore, dissection under the hilar plate is difficult and unsafe. The elements of the segmental or

sectional pedicle must supply the segment or section into which the pedicles enter. There is no need to consider any injury to the elements that supply the remaining liver segments when the pedicles above the hilar plate are interrupted.

Ramifications of the Glissonean pedicle and liver anatomical classification

The classification of liver anatomy and resections was defined in the International Hepato-Pancreatico-Biliary Association (IHPBA) Brisbane 2000 terminology and was reviewed by Strasberg in 2005. The second-order division of the classification in the IHPBA Brisbane 2000 terminology is based on the ramification of the hepatic artery and bile duct. On the

other hand, the classification according to the Glissonean pedicles is different from that according to the IHPBA Brisbane 2000 terminology. Specifically, the secondary Glissonean pedicles divide into 3 branches at the hepatic hilus. These pedicles can be approached extrahepatically without liver dissection. Therefore, the liver can be divided into 3 segments according to the ramification of the secondary Glissonean pedicles. This classification is called Takasaki's classification. By comparison, the left segment in Takasaki's classification corresponds to the left medial and left lateral sections in the Brisbane 2000 system. In Takasaki's segmentations, there are hepatic veins between segments.

Table 1: Comparison of classification systems of liver anatomy and resection

Healey	Couinaud	Takasaki	Brisbane 2000 system
Right lobe	Right liver		Right liver
Posterior segment (S6S7)	Right lateral sector (S6S7)	Right segment (S6S7)	Right posterior section (S6S7)
Anterior segment (S5S8)	Right paramedian sector (S5S8)	Middle segment (S5S8)	Right anterior section (S5S8)
Left lobe	Left liver		Left liver
Medial segment (S4)	Left paramedian sector (S3S4)	Left segment (S2S3S4)	Left medial section (S4)
Lateral segment (S2S3)	Left lateral sector (S2)		Left lateral section (S2S3)
Caudate lobe (S1)	Caudate lobe (S1)	Caudate area (S1)	Segment 1 (S1)

Extra-Fascial approach to the secondary Glissonean pedicles at the hepatic hilus

Couinaud described three approaches to the hepatic hilus in his book entitled Surgical Anatomy of the Liver Revisited. Conventional dissection in the hepatoduodenal ligament is referred to as the intra-fascial approach. This procedure has also been referred to as a controlled hepatectomy method. In this approach, dissection is performed under the hilar plate, making this procedure unsafe, and surgeons have to consider variations in the hepatic artery and bile ducts. In the extra-fascial approach, the Glissonean pedicles can be dissected from the liver parenchyma extrahepatically at the hepatic hilus. The main portal fissure or left supra-hepatic fissure is opened after dissecting the liver parenchyma and surgeons must confirm the pedicles that arise from the hilar plate or umbilical plate. This procedure is referred to as the extra-fascial and transfissural approach introduced by Lin and Tung. The left pedicle can be approached at the left end of the hilar plate. This approach was described by Couinaud in 1985. In brief, the hilar plate is detached from the liver parenchyma. Then, the ligamentum venosum is confirmed and the left pedicle is dissected along the left pedicle at the ventral side of the ligamentum venosum. The lesser omentum is not necessarily divided. The tertiary branch of the caudate lobe and the right posterior bile duct branch from the left hepatic duct at the dorsal side of ligamentum venosum. Therefore, surgeons do not have to consider any variations of the hepatic bile duct. After dissection around the pedicle, the pedicle is encircled and clamped. The left pedicle is divided following the same procedure of the right side pedicle transection.

However, the left pedicle is the thickest of the pedicles and therefore requires careful transection. A stapler is occasionally useful for transecting the left pedicle.

Approach the tertiary branches

One secondary Glissonean pedicle has 6–8 tertiary branches. The territory of a single tertiary branch does not correspond to Couinaud's segment. We therefore refer to the area fed by 1 tertiary branch as a cone unit of the liver. Couinaud's segments consist of 3 or 4 cone units. The tertiary branches in the left liver can easily be approached because they can be observed along the umbilical portion without liver dissection. However, in the right liver, the tertiary branches cannot be observed at the hepatic hilus. We should therefore clamp the pedicle of the right anterior section at the hepatic hilus and confirm the border between the sections. The tertiary branches close to the hepatic hilus can be approached around the secondary pedicle near the hepatic hilus. However, the tertiary branches that originate from the deep portions of the secondary pedicles cannot be approached from the hepatic hilus. These deep branches can be approached after initially dissecting the liver parenchyma on the border between the sections. The border between the sections can be confirmed by clamping the secondary pedicle at the hepatic hilus. If the tertiary branch that possibly feeds the territory which includes the tumor is found, the tertiary branch is clamped and the territory is confirmed.

Our experience and results

We have performed 17 hepatic resections using this technique.

Indications were hepatocellular carcinomas and metastases from colorectal cancers with curative intent. Fifteen were major hepatic resections (12 right hepatectomies and one right trisegmentectomy and one left hepatectomy all for hepatocellular carcinoma) and two segmental resections (one segment 6 ANF one segment 3 both for colo-rectal metastases). Procedure was evaluated with respect to the feasibility, duration of surgery and amount of blood loss.

In our study the average blood loss for major resection was 500ml and it was 200ml for segmental resections. Average duration of surgery was 4 hours for major and 2 hours for segmental resections.




Fig 1: Anatomy of Glissonean hepatic pedicles

Discussion

Anatomical hepatectomy has been performed more easily using the Glissonean approach than by using controlled hepatectomy methods, which are exemplified by the intra-fascial approach of Couinaud. In particular, small anatomical hepatectomies such as sectionectomy or Couinaud's segmentectomy are difficult to perform using controlled hepatectomy methods. When surgeons plan to carry out anterior sectionectomy using controlled hepatectomy methods, it takes a long time to dissect and confirm each branch of the hepatic artery, the portal vein and the bile duct to the anterior section. False confirmation of the hepatic artery or bile duct occasionally occurs. Makuuchi, *et al* indicated anterior sectionectomy as the most difficult type of hepatic resection. However, surgeons can determine the area of the anterior section when one of the secondary Glissonean pedicles to the anterior section is dissected and clamped using the Glissonean pedicle approach. This procedure will shorten the operation time and reduce the stress on surgeons because they do not have to consider any anomalies of the arteries and bile ducts using this procedure. Anatomical hepatectomy is different from partial hepatectomy, which does not consider the area that is fed by the sectional or segmental portal pedicle. In anatomical hepatectomy, the surgeon first ligates the portal pedicle and then confirms the area fed by the portal pedicle. After this, the surgeon identifies the tumor location and dissects the liver parenchyma along the border between the area fed and the area not fed by the portal pedicle. This anatomical hepatectomy approach is appropriate for liver tumors that invade the peripheral portal tract such as hepatocellular carcinoma (HCC). Because liver tumors invade the portal vein, tumor cells are disseminated to the sinusoids around the tumor and spread through the area which is fed by the portal vein. Therefore,

anatomical hepatectomy is indicated for HCC. The advantages of anatomical hepatectomy for HCC have been indicated in previous studies. The advantage of the Glissonean pedicle approach is that it is a simple method which can separate the liver into 3 sections by simply clamping the secondary Glissonean pedicle extrahepatically without liver dissection at the hepatic hilus. The main hepatic vein, either the right hepatic vein or the middle hepatic vein, is present on the border between the sections. The relationship between the area of the section and the tumor location can be easily noted preoperatively by the Glissonean pedicle approach. Sectionectomy is considered as fundamental anatomical hepatectomy; however, when liver function is insufficient for sectionectomy, changing to smaller anatomical resections, such as cone unit resection, can be achieved by using the tertiary branches. Therefore, liver surgeons should know the Glissonean pedicle approach and should be able to perform any type of anatomical hepatectomy.




Fig 2: Isolation of Right Posterior Glissonean Pedicle

References

1. Lortat-Jacob JL, Robert HG. Hepatectomie Droite Reglee. *Presse Med.* 1952;60:549–551.
2. Lin TY, Chem KM, Liu TK. Total right hepatic lobectomy for primary hepatoma. *Surgery.* 1960;48:1048–1060.
3. Tung TT, Quang ND. A new technique for operating on the liver. *Lancet.* 1963;26:192–193. DOI: 10.1016/S0140-6736(63)91210-8.
4. Takasaki K, Kobayashi S, Tanaka S, Muto H, Watayo T, Saito A, *et al.* Newly developed systematized hepatectomy by Glissonean pedicle transection method *Shujutsu.* 1986;40:7–14.
5. Launois B, Jamieson GG. The importance of Glisson's capsule and its sheath in the intrahepatic approach to resection of the liver. *Surg Gynecol Obstet.* 1992;174:7–10.
6. Strasberg SM. Nomenclature of hepatic anatomy and resections: A review of the Brisbane 2000 system. *J Hepatobiliary Pancreat Surg.* 2005;12:351–355. DOI: 10.1007/s00534-005-0999-7.
7. Takasaki K. Glissonean pedicle transection method for hepatic resection: a new concept of liver segmentation, *J Hepatobiliary Pancreat Surg.* 1998;5:286–291. DOI: 10.1007/s005340050047.