



E-ISSN: 2616-3470

P-ISSN: 2616-3462

© Surgery Science

www.surgeryscience.com

2022; 6(1): 18-20

Received: 13-11-2021

Accepted: 15-12-2021

Surya Rao Rao Venkata Mahipathy
Professor & Head, Department of
Plastic & Reconstructive Surgery,
Saveetha Medical College &
Hospital, Thandalam,
Kanchipuram, Tamil Nadu, India

Alagar Raja Durairaj
Professor, Department of Plastic &
Reconstructive Surgery, Saveetha
Medical College & Hospital,
Thandalam, Kanchipuram,
Tamil Nadu, India

Narayanamurthy Sundaramurthy
Associate Professor, Department of
Plastic & Reconstructive Surgery,
Saveetha Medical College &
Hospital, Thandalam,
Kanchipuram, Tamil Nadu, India

Anand Prasath Jayachandiran
Assistant Professor & Head,
Department of Plastic &
Reconstructive Surgery, Saveetha
Medical College & Hospital,
Thandalam, Kanchipuram,
Tamil Nadu, India

Suresh Rajendran
Senior Resident, Department of
Plastic & Reconstructive Surgery,
Saveetha Medical College &
Hospital, Thandalam,
Kanchipuram, Tamil Nadu, India

Corresponding Author:

Surya Rao Rao Venkata Mahipathy
Professor & Head, Department of
Plastic & Reconstructive Surgery,
Saveetha Medical College &
Hospital, Thandalam,
Kanchipuram, Tamil Nadu, India

Pediced anterolateral thigh flap cover for a post-infective raw area of the inguinal region following a femoral artery graft: A case report

Surya Rao Rao Venkata Mahipathy, Alagar Raja Durairaj, Narayanamurthy Sundaramurthy, Anand Prasath Jayachandiran and Suresh Rajendran

DOI: <https://doi.org/10.33545/surgery.2022.v6.i1a.811>

Abstract

Pediced anterolateral thigh (ALT) flaps are useful flaps for ipsilateral groin defects without the difficulty of microvascular surgery. Pediced ALT flap has many advantages over other regional flaps like having a large skin and soft tissue availability, the good pedicle length, wide arc of rotation and being reliable. The pediced anterolateral thigh flap is a versatile option for coverage of soft tissue defects within its reach. Here, we report a case of a groin defect which was a post-infective sequelae following a vascular graft for the femoral artery for peripheral vascular disease which was resurfaced with a pediced anterolateral thigh flap from the ipsilateral side.

Keywords: groin defect, pediced, anterolateral thigh flap, no complication, versatile

Introduction

Anterolateral thigh (ALT) flap is a popular workhorse flap for reconstruction of soft tissue of various regions, first described by Song *et al.* [1]. The versatility of this flap is due to its long vascular pedicle, large size of skin paddle, ability to harvest different tissues in composite nature, its use as a sensate and flow through flap and minimal to no donor site morbidity. The two team approach has decreased the operative time and thereby decreasing the anaesthetic complications significantly [2]. The pediced ALT flap is also versatile due to the similar features [3]. The vascular supply to the skin paddle of ALT flap is by septocutaneous or musculocutaneous perforators from the descending branch of lateral circumflex femoral artery. The descending branch anastomoses to the lateral superior genicular artery or the profunda femoris artery perforators thereby allowing the ALT flap to be harvested based on either its proximal or distal circulation. The proximal based circulation along with the long pedicle can be utilised to cover the various soft tissue defects involving the abdomen, groin, perineum, and trochanteric regions and soft tissue defects around the knee as a distally based circulation [3].

Case Report

53 year old male presented to us with a raw area of the right groin since 3 weeks. It started as a small ulcer and has progressed to the present size. He was a known case of peripheral vascular disease for he underwent left above knee amputation 3 years ago. Now, he has right femoral thrombosis for which it was grafted by the vascular surgeon team. But he developed wound infection and dehiscence of the suture line and wound with exposure of the vascular graft. He was a known diabetic for which he was on insulin and oral hypoglycaemic agents. On examination, there was an unhealthy area right groin with suture line dehiscence and a minimal raw area. We planned for debridement of the wound and cover with a pediced anterolateral thigh flap was marked on the ipsilateral side. (Fig. 1) Under epispinal anaesthesia, thorough debridement was done and the pediced ALT flap and the perforator from the lateral circumflex femoral artery was identified and marked. (Fig. 2) Flap was elevated and pedicle dissection was done. Flap was inset into the defect with 2-0 nylon sutures and the secondary defect was also closed primarily. (Fig. 3) Dressing was done after achieving haemostasis. Post-operative period

was uneventful with the flap well settled and sutures were removed on the 14th post-operative day. (Fig. 4)



Fig 1: Area to be debrided and the anterolateral thigh flap marked



Fig 2: Defect after debridement



Fig 3: Immediate post-operative picture showing the pedicled ALT flap



Fig 4: Late post-operative picture showing a well settled flap

Discussion

Pedicled ALT flap is a versatile option for reconstruction of soft

tissue defects around the groin region^[4]. The rotation arcs of the ALT flap include the umbilicus superiorly, ipsilateral posterior superior iliac spine (PSIS) laterally, and tibial tuberosity below^[5]. Pedicled ALT flap can also be used to cover the hand soft tissue defects of the hand. Lannol *et al.* had the largest series of the proximally pedicled ALT for abdominal and pelvic reconstruction where they described the farthest reach of the ALT flap to the lower costal margin, posterior superior iliac spine, anterior margin of anus, and contralateral iliac fossa^[6]. Wang *et al.* described pedicled ALT flap as a reliable flap in the management of recurrent trochanteric pressure sores.⁷ The pedicled ALT flap was also used to resurface ischial pressure ulcers by Lee JT *et al.*^[8] Lee GK *et al.* emphasized the pedicled ALT as the preferred method for reconstruction of total penile defects. Pedicled ALT flap had several advantages over the free radial forearm flap for total penile reconstruction like lack of need of microvascular surgery, better colour match, bulky flap, and less donor site morbidity as mentioned by Lee GK *et al.*^[9] Many authors have described the pedicled ALT flap for pelvic defects^[10-12] Zelken JA *et al.* utilised pedicled ALT flap along with groin flap to reconstruct the mangled hand injuries^[13]. ALT flap is a versatile option to cover the soft tissue defects around the knee joint and upper leg based on the distal circulation, but venous congestion is a problem which can be prevented by supercharging the flap vein to great saphenous vein^[14, 15]. Bulkiness of the flap was not a problem in our series. ALT flap was thin and pliable in most of the non-obese patients we encountered. All patients were satisfied with the contour of the flap postoperatively. Bharath *et al* have used the pedicled ALT flap to resurface contralateral groin defects^[16]. Ahmed *et al.* described a series of 17 pedicled ALT flaps, tunneling of the flap between the rectus and the fascial septa for ipsilateral inguinal defects^[17]. Complex defects of the groin following cancer ablation can be large and difficult to resurface, but was easily achieved by the pedicled ALT flaps^[18]. The pedicled anterolateral thigh flap is therefore a versatile, thin, pliable, flap useful for complex reconstructions^[19, 20].

Conclusion

The pedicled ALT flap is a reliable, safe and versatile flap used to cover complex defects of the groin region by its wide the arc of rotation and long pedicle length and minimal donor site morbidity. It can be used to cover the defects of the inguinal, iliac and pelvic and abdominal wall defects.

References

1. Song YG, Chen GZ, Song YL. The free thigh flap: A new free flap concept based on the septocutaneous artery. *British Journal of Plastic Surgery.* 1984;37:149-159.
2. Wei FC, Jain V, Celik N, Chen HC, Chuang DC, Lin CH. Have we found an ideal soft-tissue flap? An experience with 672 anterolateral thigh flaps. *Plastic and Reconstructive Surgery.* 2002;109:2219-2226. discussion 2227-2230.
3. Gravvanis AI, Tsoutsos DA, Karakitsos D *et al.* Application of the pedicled anterolateral thigh flap to defects from the pelvis to the knee. *Microsurgery.* 2006;26:432-438.
4. Friji MT, Suri MP, Shankhdhar VK, Ahmad QG, Yadav PS. Pedicled anterolateral thigh flap: a versatile flap for difficult regional soft tissue reconstruction. *Annals of Plastic Surgery.* 2010;64:458-461.
5. Ravikiran NS, Chauhan S, Bhattacharyya MS. The Versatility of Pedicled Anterolateral Thigh Flap: A Tertiary Referral Center Experience from India. *International Microsurgery Journal.* 2017;1(2):5.

6. Lannon DA, Ross GL, Addison PD, Novak CB, Lipa JE, Neligan PC. Versatility of the proximally pedicled anterolateral thigh flap and its use in complex abdominal and pelvic reconstruction. *Plastic and Reconstructive Surgery*. 2011;127:677-688.
7. Wang CH, Chen SY, Fu JP, *et al*. Reconstruction of trochanteric pressure sores with pedicled anterolateral thigh myocutaneous flaps. *Journal of Plastic, Reconstructive & Aesthetic Surgery*. 2011;64:671-676.
8. Lee JT, Cheng LF, Lin CM, Wang CH, Huang CC, Chien SH. A new technique of transferring island pedicled anterolateral thigh and vastus lateralis myocutaneous flaps for reconstruction of recurrent ischial pressure sores. *Journal of Plastic, Reconstructive & Aesthetic Surgery*. 2007;60:1060-1066.
9. Lee GK, Lim AF, Bird ET. A novel single-flap technique for total penile reconstruction: the pedicled anterolateral thigh flap. *Plastic and Reconstructive Surgery Z*. 2009;124:163-166.
10. Gentileschi S, Servillo M, Garganese G, Simona F, Scambia G, Salgarello M. Versatility of pedicled anterolateral thigh flap in gynecologic reconstruction after vulvar cancer extirpative surgery. *Microsurgery*. 2017;37:516-524.
11. Zelken JA, AlDeek NF, Hsu CC, Chang NJ, Lin CH, Lin CH. Algorithmic approach to lower abdominal, perineal, and groin reconstruction using anterolateral thigh flaps. *Microsurgery*. 2016;36:104-114.
12. Wong S, Garvey P, Skibber J, Yu P. Reconstruction of pelvic exenteration defects with anterolateral thigh-vastus lateralis muscle flaps. *Plastic and Reconstructive Surgery*. 2009;124:1177-1185.
13. Zelken JA, Chang NJ, Wei FC, Lin CH. The combined ALT-groin flap for the mutilated and degloved hand. *Injury*. 2015;46:1591-1596.
14. Pan SC, Yu JC, Shieh SJ, Lee JW, Huang BM, Chiu HY. Distally based anterolateral thigh flap: an anatomic and clinical study. *Plastic and Reconstructive Surgery*. 2004;114:1768-1775.
15. Lin CH, Zelken J, Hsu CC, Lin CH, Wei FC. The distally based, venous supercharged anterolateral thigh flap. *Microsurgery*. 2016;36:20-28.
16. Bharath SP, Madhusudan G, Manjunath S. Pedicled anterolateral thigh flap for contralateral groin composite defect. *Indian Journal of Plastic Surgery*. 2010;43:103-105.
17. Ahmad QG, Reddy M, Shetty KP, Prasad R, Hosi JS, Bhathena M. Groin reconstruction by anterolateral thigh flap: A review of 16 cases. *Indian J Plast Surg*. 2004;37:34-9.
18. Evriviades D, Raurell A, Perks AG. Pedicled anterolateral thigh flap for reconstruction after radical groin dissection. *Urology*. 2007;70:996-9.
19. Kimura N, Satoh K. Consideration of a thin flap as an entity and clinical applications of the thin anterolateral thigh flap. *Plastic and Reconstructive Surgery*. 1996;97(5):985-992.
20. Park JE, Rodriguez ED, Bluebond-Langer R *et al*. The anterolateral thigh flap is highly effective for reconstruction of complex lower extremity trauma. *The Journal of Trauma*. 2007;62(1):162-165.