



E-ISSN: 2616-3470

P-ISSN: 2616-3462

© Surgery Science

www.surgeryscience.com

2022; 6(4): 01-04

Received: 01-07-2022

Accepted: 05-08-2022

**Dr. Khan Tauqir Ahmad Khan
Ajmal Khan**

Assistant Professor, Department of
General Surgery, LTMMC Sion,
Maharashtra, India

Dr. Vahid Saleem Shaikh

Assistant Professor, Department of
General Surgery, Dr. VM Medical
College, Solapur, Maharashtra,
India

Dr. Sandip Dadmal

Assistant Professor, Department of
General Surgery, Dr. VM Medical
College, Solapur, Maharashtra,
India

Clinical study and management of fistula in Ano

**Dr. Khan Tauqir Ahmad Khan Ajmal Khan, Dr. Vahid Saleem Shaikh and
Dr. Sandip Dadmal**

DOI: <https://doi.org/10.33545/surgery.2022.v6.i4a.939>

Abstract

Background: Fistula in Ano is one disease which is easy to diagnose but difficult to cure as patients report in very late phase of the disease and it leads to problems. Aim of this study was to determine the etiology, clinical presentation, efficacy of different surgical approach, early and late post-operative complications, and recurrence rate in cases of fistula in Ano.

Methods: Prospective Observational Study was conducted in Department of General Surgery, Government Medical College, Solapur from Sep 2016 to Aug 2018 with Sample size of 50 cases.

Results: Most common age group affected between 18-29 years, with male preponderance. Most common etiological factor is perianal sepsis following Crypto glandular infection, with pain as most prevalent symptom. Simple fistula is more common, and in complex fistula most prevalent type is high transsphincteric fistula. Most common procedure done is fistulotomy. Most common complication is recurrence. Most number of complications was seen in fistulotomy. Maximum recurrence is seen after fistulotomy.

Conclusion: Most common cause is crypto glandular infection, most common feature is pain, Fistulectomy is more efficacious, and complications are maximum with fistulotomy, highest recurrence seen after fistulotomy.

Keywords: Fistula, pain, setn, fistulotomy, Fistulectomy, set on, recurrence

Introduction

Anal Fistula-in-Ano is a chronic abnormal communication between the epithelized surface of anal canal or rectum and usually the perianal skin ^[1]. There are many causes of anorectal suppuration, the most common being the infection of Crypto glandular structures ^[2]. If the right kind of surgery is not performed, the disease tends to recur. It can also recur due to negligence in post-operative care ^[3]. The prevalence of nonspecific anal fistulae has been estimated to be 8.6 to 10/100,000 of the population per year, with a male to female ratio of 1.8:1 ^[4]. Park classified fistula in Ano into four different types depending on its relation to the anal sphincters – Intersphincteric – 70%, Transsphincteric - 25%, Suprasphincteric - 5%, Extrasphincteric - 1% ^[5]. Goodsall's rule predicts that if a transverse anal line is drawn transversely through the anus, an external opening anterior to this transverse anal line will have a radial tract to open in the dentate line, whereas an external opening posterior to the transverse anal line will have a curved fistulous tract to open in the posterior midline. The exception to this rule is an anterior fistula with its external opening lying 3 cm or more from the anal verge will have a curved path like a posterior fistula to have its internal opening in the posterior midline ^[6]. A fistulotomy is the surgical opening of a fistulous tract ^[7]. Fistulotomy is the procedure of choice for simple intersphincteric fistulous tract and also for low lying transsphincteric fistulae ^[8]. Seton fistulotomy is the procedure of choice for high anal fistulas involving significant amount of external anal sphincter ^[9, 10].

Fistulectomy is a procedure of excising the fistulous tract from external opening up to the internal opening ^[11].

Fistulectomy removes secondary fistulous tracts compared to fistulotomy, but fistulotomy has shorter healing times and less chance of damage to the sphincters ^[12]. Many new techniques such as the mucosal advancement flap, the Gore Bio-A fistula plug, and the ligation of intersphincteric fistulous tract procedure have seen recent acceptance ^[13].

Corresponding Author:

**Dr. Khan Tauqir Ahmad Khan
Ajmal Khan**

Assistant Professor, Department of
General Surgery, LTMMC Sion,
Maharashtra, India

Aims & Objective

1. To study the etiology of fistula in Ano.
2. To study the clinical presentation of patients having fistula in Ano.
3. To study the efficacy of different surgical approach i.e. fistulotomy,
4. Fistulectomy, and set on treatment in management of fistula in Ano.
5. To study early and late post-operative complications in these patients.
6. To study the recurrence rate in these patients.

Materials and Methods

Type of study: Prospective Observational Study.

Place of study: Department of General Surgery, Government Medical College, Solapur.

Period of study: Duration from Sep 2016 to Aug 2018.

Sample size: 50 cases.

After admission detailed history was taken to establish proper diagnosis. Thorough physical examination was done in each case. Baseline investigations, as routinely required were done followed by imaging studies. After proper evaluation and preparation, patients who required surgical management were taken up for surgery. Meticulous techniques were practiced.

Results**Table 1:** Distribution of patients according to age in years

| Sr. No | Age group in years | Number of cases (%) | Mean age of patients |
|--------|--------------------|---------------------|----------------------|
| 1 | 18 - 29 | 19 (38%) | 22.84 |
| 2 | 30 - 39 | 17 (34%) | 34.76 |
| 3 | 40 - 49 | 06 (12%) | 44.50 |
| 4 | 50 - 59 | 05 (10%) | 55.80 |
| 5 | > 60 | 03(06%) | 69.00 |

Maximum 38% patients are from age group 18-29 years and minimum 06% patients are from age group > 60 years.

Table 2: Distribution of Patients according to gender

| Sr. No | Sex/Gender | Number of cases (%) |
|--------|------------|---------------------|
| 1 | Male | 37 (74%) |
| 2 | Female | 13 (26%) |
| | Total | 50 (100%) |

The distribution of patients according to gender show 74% patients are male and 26% patients are female.

Table 3: Etiology/predisposing factors of fistula in Ano

| Sr. No | Etiology/Predisposing factors | Number of cases (%) |
|--------|--|---------------------|
| 1 | Perianal Sepsis/Crypto glandular infection | 31 (62%) |
| 2 | Tuberculosis | 03 (06%) |
| 3 | Malignancy | 00 (00%) |
| 4 | Fissure in ano | 04 (08%) |
| 5 | Crohn's Disease | 00 (00%) |
| 6 | No previous significant history | 10 (20%) |
| 7 | Trauma in perianal region | 02 (04%) |
| | Total | 50 (100%) |

In present study most common etiological factor is perianal sepsis following Crypto glandular infection 62% while no previous significant history is found in 20% cases.

Table 4: Clinical presentation of patients

| Sr. No. | Symptoms | Number of patients | Percentage |
|---------|--------------|--------------------|------------|
| 1 | Pain | 16 | 32% |
| 2 | Discharge | 15 | 30% |
| 3 | Swelling | 09 | 18% |
| 4 | Itching | 04 | 08% |
| 5 | Constipation | 03 | 06% |
| 6 | Bleeding | 01 | 02% |
| 7 | Fever | 02 | 04% |

Most prevalent symptom is pain in 32%. Bleeding is the rarest symptoms present in only 02% patient.

Table 5: Classification of Fistula

| Sr. No. | Type of fistula | Number of patients | Percentage |
|---------|-----------------|--------------------|------------|
| 1 | Simple fistula | 38 | 76% |
| 2 | Complex fistula | 12 | 24% |

It is found that simple fistula is more common 76% cases than complex fistula 24% cases.

Table 6: Type of complex Fistula

| Sr. No. | Type of complex fistula | Number of patients | Percentage |
|---------|-------------------------|--------------------|------------|
| 1 | High transsphincteric | 07 | 58.33% |
| 2 | Suprasphincteric | 00 | 00% |
| 3 | Extrasphincteric | 00 | 00% |
| 4 | Multiple openings | 03 | 25% |
| 5 | Recurrent | 02 | 16.67% |

Among complex fistula most common is high transsphincteric fistula 58.33% cases

Table 7: Distribution of patients according to operative procedures

| Sr. No. | Operative Procedure | Number of cases | Percentage |
|---------|---------------------|-----------------|------------|
| 1 | Fistulectomy | 17 | 34% |
| 2 | Fistulotomy | 28 | 56% |
| 3 | Seton | 05 | 10% |
| | Total | 50 | 100% |

In present study most common procedure done is fistulotomy 56% patients. Seton is done in only in 10% patients.

Table 8: Postoperative complications

| Sr. No. | Complications | Number of cases | Percentage |
|---------|-------------------|-----------------|------------|
| 1 | Persisting Sepsis | 02 | 04% |
| 2 | Incontinence | 01 | 02% |
| 3 | Recurrence | 13 | 26% |
| | Total | 16 | 32% |

It is found that complications are seen in 16 (32%) cases out of 50 fistula in Ano surgeries, out of which most common complication is recurrence 26% cases.

Table 9: Analysis of complications with respect to procedure

| Sr. No. | Complications | Fistulectomy | Fistulotomy | Seton | Total |
|---------|-------------------|--------------|-------------|-------|-------|
| 1 | Persisting Sepsis | 00 | 01 | 01 | 02 |
| 2 | Incontinence | 00 | 01 | 00 | 01 |
| 3 | Recurrence | 03 | 10 | 00 | 13 |
| | Total | 03 | 12 | 01 | 16 |

The Chi-square statistic with Yates correction is 1.5415.

The p –value is 0.21439. Not significant at $p < 0.05$

Most number of complications are present in fistulotomy 12 out of 28 i.e. (42.85%) cases, and much less in case of Fistulectomy only 3 out of 17 i.e. (17.64%). The difference was found to be statistically not significant as p- value is greater than 0.05.

Table 10: Recurrence rate in various surgical procedures for Fistula in Ano

| Sr. No. | Surgical Procedures | Number of cases | Recurrence |
|---------|---------------------|-----------------|-------------|
| 1 | Fistulectomy | 17 | 03 (17.64) |
| 2 | Fistulotomy | 28 | 10 (35.71%) |
| 3 | Seton | 05 | 00 (00%) |

The Chi-square statistic with Yates correction is 0.1758.

The p – value is 0.674971. Not significant at $p < 0.05$

Maximum recurrence is seen after fistulotomy 35.71% cases, with no recurrence after Seton procedure.

The difference was found to be statistically not significant as p value is greater than 0.05.

Discussion

Maximum i.e. 38% patients are from age group 18-29 years, in contrast Vyas A. K *et al.* [14] observed most common age group was 22 to 55 years. In present study, male patient is 74%, i.e. >50%, similar result were observed studies by Tated S.P *et al.* [3] 82.71%, Alexander N *et al.* [13] 82.5%, Jebakumar A *et al.* [4] 80%. Most common etiological factor is perianal sepsis following Crypto glandular infection 62%, similar result was observed by Alexander N *et al.* [13]. In present study most prevalent symptom is pain 32%, in contrast Tated S.P *et al.* [3] 97.53% presented with swelling in perineal region, Sushma R *et al.* [15] 96% presented with perianal discharge. We found that simple fistula (76%) is more common, similar result observed by Alexander N *et al.* [13] 85%.

In case of complex fistula, most numbers are of high transsphincteric i.e. 58.33%, similar result observed by Alexander N *et al.* [13].

In present study most common procedure done is fistulotomy 56%, in contrast Tated S.P *et al.* [3] 74.07% patients were treated by Fistulectomy, Jebakumar A *et al.* [4] Fistulectomy surgery had done in 68%.

It is found that complications are seen in 32% cases of fistula in Ano surgery, out of which most common complication is recurrence 26%. Most number of complications are present in fistulotomy 42.85% cases, in contrast to present study Ramachandra M. L *et al.* [16] recurrence seen in 25% cases of Fistulectomy and 10% in fistulotomy. Kharadi A *et al.* [17] fistulotomy has higher incidence of recurrence than Fistulectomy as more tissue is left behind. RCT by Kronberg, the recurrence rate following fistulotomy and Fistulectomy were reported to be 12.5% and 09.5% respectively [18]. In this study, it is found that maximum recurrence seen after fistulotomy 35.71%) cases, in contrast to present study, Vyas AK *et al.* [14], showed that there was 3.2% in Fistulotomy.

History

Anal fistula has been described virtually from beginning of medical history. Hippocrates in about 430 B.C., suggested that the disease was caused by “contusions and tubercles occasioned by rowing or riding on horseback”. Ayurvedically medicated set on, “kshara” used by Sushruta for treating fistula in Ano.

Conclusion

Commonest etiological factor associated with Fistula in Ano is chronic nonspecific inflammation most commonly due to Crypto glandular infection, with pain as most common presentation. Fistulectomy is more efficacious than Seton and Fistulotomy, in our study. Post-operative complications are maximum with fistulotomy. Highest recurrence rate is observed following Fistulotomy and no recurrence seen with Seton.

Conflict of Interest

Not available

Financial Support

Not available

References

1. Merck Manual Consumer Version. Retrieved 2016 Jun 27.
2. Yoon S. Fischer's Mastery of Surgery. Annals of Surgery. 2012 Oct 1;256(4):669.
3. Tated SP, Sharma K, Hatkar AA. Clinical study of various modalities of treatment for fistula in Ano at a tertiary care hospital. International Surgery Journal. 2017;4(11):3670-4.
4. Jebakumar A, Shanmugam S, Anbarasan. A comprehensive study of fistula in Ano. IAIM, 2016;3(8):66-72.
5. Parks AG, Gordon PH, Hardcastle JD. A classification of fistula-in-Ano. British Journal of Surgery. 1976 Jan;63(1):1-2.
6. Townsend CM, Beauchamp RD, Evers BM, Mattox KL. Sabiston Textbook of Surgery E-Book. Elsevier Health Sciences; c2016 Apr 22.
7. Jain BK, Vaibhaw K, Garg PK, Gupta S, Mohanty D. Comparison of a fistulectomy and a fistulotomy with marsupialization in the management of a simple anal fistula: a randomized, controlled pilot trial. Journal of the Korean Society of Coloproctology. 2012 Apr;28(2):78.
8. Corfitsen MT, Hansen CP, Christensen TH, Kaae HH. Anorectal abscesses in immunosuppressed patients. The European journal of surgery Acta chirurgica. 1992;158(1):51-3.
9. Deen KI, Williams JG, Hutchinson R, Keighley MR, Kumar D. Fistulas in Ano: endoanal ultrasonography assessment assists decision making for surgery. Gut. 1994;35(3):391-4.
10. Garcia-Aguilar J, Belmonte C, Wong W, Goldberg DW, Madoff RD. Cutting set on versus two-stage set on fistulotomy in the surgical management of high anal fistula. British Journal of Surgery. 1998;85(2):243-5.
11. Del Pino A, Nelson RL, Pearl RK, Abcarian H. Island flap anoplasty for treatment of transsphincteric fistula-in-Ano. Diseases of the colon & rectum. 1996;39(2):224-6.
12. GH RG, JL MV. Fistulotomy vs Fistulectomy. Ultrasonography evaluation of lesion of the anal sphincter function. Revisit de gastroenterological de Mexico. 1999 Oct 1;64(4):167-70.
13. Alexander N, Sanniyasi S, Joseph J, Paramasivam S. A Single Center Study of 80 Cases of Fistula-in-Ano. International Journal of Scientific Study. 2016;4(8):122-5.

14. Vyas AK, Katlana A, Singh A, Thakur S, Yadav A. Incidence of Low Fistula in Ano and Results of Fistulotomy in Rural Tertiary Centre. *Ann Int Med Den Res.* 2017;3:3.
15. Ramteke S, Gupta AM, Soni P. Clinical study of fistula in Ano. *Journal of Evidence Based Medicine and Healthcare.* 2017;4(11):595-9.
16. Ramachandra ML, Garg M. A comparative study in the management of fistula in Ano using various modalities. *International Surgery Journal.* 2018;5(6):2223-7.
17. Kharadi A, Varikoo V. A descriptive analysis of management of fistula in-Ano. *International Surgery Journal.* 2016;3(2):683-6.
18. Kronberg O. To lay open or excise fistula-in-Ano: A randomized trial. *Br J Surgery.* 1985;72(1):970.

How to Cite This Article

Khan KTAKA, Shaikh VS, Dadmal S. Clinical study and management of fistula in Ano. *International Journal of Surgery Science.* 2022;6(4):01-03.
DOI: <https://doi.org/10.33545/surgery.2022.v6.i4a.939>

Creative Commons (CC) License

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.